

Research Article

## Studies on Biochemical Parameters of Probiotic Supplemented Diet on Rohu (*Labeo rohita*) Fresh Water Fish Fingerlings

J. Gopinath\*, R. Ramanibai, V. Krishnakumar and S. Inbasekaran

Department of Zoology, Unit of Aquatic Biodiversity, University of Madras, Guindy Campus,  
Chennai-600 025, Tamil Nadu, India  
gopije@gmail.com\*

Received: March 01 2019/Accepted: 30 April 2019/Published: 07 May 2019

### Abstract

Microbial diseases are one of the major problems in fresh water aquaculture. There is an urgent need to develop microbial control strategies since disease epidemics are recognized as important constraints to aquaculture production. This study reports the effect of the gram positive probiotic bacteria *Lactobacillus* sp. on the freshwater fish Rohu (*Labeo rohita*) fingerlings against a pathogenic bacteria *Vibrio cholerae* of first feeding. Probiotic was administered through the diet at four different concentrations such as  $312 \times 10^{-4}$  (Group I),  $126 \times 10^{-5}$  (Group II),  $75 \times 10^{-6}$  (Group III) and  $22 \times 10^{-7}$  (Group IV) cfu/g of feed and fed at the rate of 3% body weight for a period of 30 d. Control group was fed without probiotic incorporated diet. The growth was estimated by morphometric measurements for healthy and disease free fingerlings, average body weight of  $22.5 \pm 1.5$  g were selected for biochemical parameters. The results indicated that protein, lipid, carbohydrate contents were high in Group III fed fish than the control group. The findings also clearly indicated the importance of probiotic feed to control *Vibrio cholerae* in freshwater aquaculture farms.

**Keywords:** *Labeo rohita*, *Vibrio cholerae*, probiotics, fingerlings, biochemical parameters.

### Introduction

Probiotics are live microbial supplements when administered in adequate quantities confer a beneficial effect on the health of the host by improving its intestinal microbial balance (Fuller, 1989; FAO/WHO, 2001). The wide use of antibiotics and chemotherapeutics to prevent and control bacterial diseases in fish farms has led to some problems like drug resistance (Alderman and Hastings, 1998; Teuber, 2001). These antibiotics may cause an inhibition of beneficial microbiota which is normally present in the digestive tract of the fish (Sugita et al., 1991). Moreover, vaccines cannot be used alone as a universal disease control measure in aquaculture (Amábile-Cuevas et al., 1995) due to their limited availability in few countries and their pathogen specific protective action for certain specific bacterial and viral diseases (FAO, 2006). Hence, the use of probiotics as biological control agents or as dietary live microbial supplements in commercial fish culture to improve growth and immune function of the fish (Gatesoupe, 1999; Irianto and Austin, 2002; Kesarcodi-Watson et al., 2008). Fish diseases are one of the major problems in the fish farm industry. Even though vaccines are being developed and marketed, they cannot be used as a universal disease control measure in aquaculture.

The use of antibiotics to cure bacterial infection and prevent fish mortality in aquaculture is becoming limited as pathogens develop resistance to the drugs (Gonzalez et al., 2000; Gomez-Gil et al., 2000). Further, beneficial bacterial flora are killed or inhibited by orally administered antibiotics, leading to efforts to find alternative disease prevention methods such as the use of non-pathogenic bacteria as probiotic bio-control agents. The use of commercial probiotics in fish is relatively ineffective as most commercial preparations are based on strains isolated from non-fish sources that are unable to survive or remain viable at high cell density in the intestinal environment of fish during the active growth phase of the fish (Gram et al., 2001). Hence, there is elegant logic in isolating putative probiotics from the host in which the probiotics intended for use. Such strains should perform better because they have already adhered to the gut wall of the fish and thus are well-adapted to compete with pathogens for nutrients. Presumably, strains that develop dominant colonies in the fish intestine are good candidates for preventing the adhesion of pathogens on the gut wall. Keeping the above points in view this study reports the effect of the gram positive probiotic bacteria *Lactobacillus* sp. on the freshwater fish Rohu (*Labeo rohita*) fingerlings against a pathogenic bacteria *Vibrio cholerae*.

\*Corresponding author

## Materials and methods

**Fingerlings collection and maintenance:** Fingerlings were collected from Porur Lake, Chennai Tamil Nadu, India. The collected fingerlings were brought to the laboratory in plastic bags with oxygenated water. Collected fingerlings were acclimatized to the laboratory conditions for 10 d in disinfected 1000 L circular FRP tanks. During acclimatization, the *Labeo rohita* fingerlings were fed with formulated diet. Healthy and disease free fingerlings, weighing average body weight and length  $22 \pm 2$  g and  $6 \pm 0.3$  cm were selected for further experiments.

**Experimental groups:** Probiotic was administered through the diet at four different concentrations such as  $312 \times 10^{-4}$  (Group I),  $126 \times 10^{-5}$  (Group II),  $75 \times 10^{-6}$  (Group III) and  $22 \times 10^{-7}$  (Group IV) cfu/g of feed and fed at the rate of 3% body weight for a period of 30 d. Control group was fed without probiotic incorporated diet.

**Biochemical analysis:** The protein content in different tissues such as gill, liver, muscle and kidney of *Labeo rohita* fingerlings was estimated according to Bradford (1976). Total lipid content in different tissues such as gill, liver, muscle and kidney of fingerlings was extracted according to the procedure of Folch et al. (1957) and estimated according to the method of Barnes and Blackstock (1973). The carbohydrate content in different tissues samples of *Labeo rohita* was estimated following the method of Roe (1955).

## Results

The efficacy of the diets was studied in different tissues such as gills, muscle, liver and kidney of *Labeo rohita* and compared with the fingerlings fed with control diet.

**Gills:** Protein content of gills in *Labeo rohita* fingerlings fed Group III was found to be  $9.68 \pm 0.27$  mg/g. However, other experimental diets showed a linear increase of protein content up to 30 d and Group III increased the protein content in gills (Fig. 1). Lipid content of gills also varied in fingerlings fed with various experimental diets. Lipid content was high in fingerlings fed with Group III ( $4.62 \pm 0.34$  mg/g). Similarly carbohydrate content was maximum in fingerlings fed with Group III ( $3.72 \pm 0.17$  mg/g). In the experimental groups, Group III diets enhanced the carbohydrate content in gills.

**Muscle:** Protein content was high in muscle of *Labeo rohita* fingerlings fed with Group III probiotic ( $21.47 \pm 0.26$  mg/g) control supplemented groups. Among the experimental groups, Group III diets showed increased protein content of the muscle. The protein content of muscle varied significantly in fingerlings fed with different experimental diets compared to the control.

Fig. 1. Protein content of gills in *Labeo rohita* fingerlings.




Fig. 2. Lipid content of liver in *Labeo rohita* fingerlings.





Fig. 3. Carbohydrate content of kidney in *Labeo rohita* fingerlings.



Lipid content was also high in muscle of fingerlings fed with Group III probiotic ( $6.44 \pm 0.27$  mg/g) and low on 30<sup>th</sup> d (Fig. 2). Among the groups, Group III diets increased the lipid content of the muscle which was significant in different experimental groups compared to the control. Carbohydrate content of muscle was high in fingerlings fed with Group III probiotic ( $5.79 \pm 0.18$  mg/g) and control supplemented diets on 30<sup>th</sup> d. Among the experimental groups, Group III diets enhanced the carbohydrate of the muscle.

The findings revealed that variations in the carbohydrate content of muscle were significant in the experimental groups compared to the control.

**Kidney:** Protein content of kidney in fingerlings fed with experimental diets varied among the groups studied. Protein content was high in fingerlings fed with Group III (11.92±0.35 mg/g) supplemented diets on 30<sup>th</sup> d. Among the experimental groups, Group III diets increased the protein content of the kidney. Lipid content of kidney in fingerlings fed with experimental diets showed significant changes among the groups studied. Lipid content was high in fingerlings fed with Group III (8.84±0.34 mg/g) on 30<sup>th</sup> d. Among the experimental groups, Group III diets enhanced the protein content of the kidney. Carbohydrate content was maximum in Group III (5.93±0.15 mg/g) on the 30<sup>th</sup> d. Among the experimental groups, Group III diets enhanced the carbohydrate content of the kidney (Fig. 3). The carbohydrate content of the kidney varied significantly in the experimental groups compared to the control.

## Discussion

Probiotic bacteria (*Lactobacillus* sp.) act against *Aeromonas* virulence in *Oreochromis mossambicus* (Who et al., 1994). In the present study, efforts were to study the effect of *V. Cholerae* in the freshwater fish *L. rohita* fed with different concentration of probiotic bacteria *Lactobacillus* sp. supplemented diets. Biochemical parameters serve as suitable indicators to assess the level of stress and iterations in metabolic cycles (Wang et al., 2005). Hence, it is necessary to understand the significance of these biochemical changes in *Labeo rohita* fingerlings, infected with *V. cholerae*. The biochemical analysis revealed that variations in the protein, lipid and carbohydrate content of gill, muscle, liver and kidney of infected fingerlings showed significant increase compared to the control and other groups. This increase may due to bacterial infection (Acha and Szyfres, 2003). Hence it is suggested that the *Lactobacillus* sp. can be used as probiotic against *V. cholerae* in the freshwater fish farming especially *Labeo rohita*. The protein content in gill, muscle, liver and kidney of *Labeo rohita* fingerlings varied in the supplemented groups compared to control. Among the different tissues studied protein content in muscle increased in fingerlings fed with probiotic supplemented diets. Lipids are essential organic constituents in animal tissue and play a key role in energy metabolism and their assessment serves as a tool for assessing the normal physiology (Palanivelu et al., 2005). Among the tissue studied, lipid content was maximum in liver tissue. The increase in the lipid content showed that the fish utilized energy derived from lipid metabolism to overcome the stress. Carbohydrate is considered to be the first organic nutrient degraded in response to stress conditions imposed on animals.

Carbohydrate content increased in gill, muscle, liver and kidney of *Labeo rohita* fingerlings fed with the probiotic Group III supplemented diets. Probiotic supplemented diet fed fingerlings showed reduction in the carbohydrate content in all tissues.

## Acknowledgements

Authors acknowledge UGC-BSR fellowship Delhi for providing financial support to carry out this study.

## References

1. Acha, P.N. and Szyfres, B. 2003. Zoonoses and communicable diseases common to man and animals. In: Bacterioses and mycoses. Pan American Health Organization (PAHO). Scientific and Technical Publication, 3rd ed Washington DC: No. 580, pp.168-179.
2. Alderman, D.J. and Hastings, T.S. 1998. Antibiotic use in aquaculture: development of antibiotic resistance-potential for consumer health risks. *Int. J. Food Sci. Tech.* 33: 139-155.
3. Amábile-Cuevas, C.F., Cardenas-Garcia, M. and Ludgar, M. 1995. Antibiotic Resistance. *Am. Sci.* 83: 320-329.
4. Barnes, H. and Blackstock, J. 1973. Estimation of lipids in fresh water animals tissues. Detailed investigation of the sulphovanilin method for total lipids. *J. Expt. Mar. Ecol.* 12: 103-118.
5. Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72: 248-254.
6. FAO. 2006. The state of world fisheries and aquaculture. Rome, FAO. p.162.
7. FAO/WHO. 2001. Evaluation of health and nutritional properties of powder milk and live lactic acid bacteria. Food and Agriculture Organization of the United Nations and World Health Organization Expert Consultation Report, Cordoba, Argentina, pp. 1-34.
8. Fernandez, R.M. 2011. Effect of Lactic acid bacteria administered orally on growth performance of *Penaeus indicus* (H. Milne Edwards) juveniles. *Res. J. Microbiol.* 6: 466-479.
9. Folch, J., Lees, M. and Stanley-Sloane, G.H. 1957. A simple method for the isolation purification of total lipids from animal tissues. *J. Biol. Chem.* 226: 497-507.
10. Gatesoupe, F.J. 1999. The use of probiotics in aquaculture. *Aquacult.* 180: 147-165.
11. Gomez-Gil, B., Rogue, A. and Turnbull, J.F. 2000. The use and selection of aquatic organisms. *Aquacult.* 191: 259-270.
12. Gonzalez, C.J., Encinas, J.P., Garcia-Lopez, M.L. and Otero, A. 2000. Characterization and identification of lactic acid bacteria from freshwater fishes. *Food Microbiol.* 17: 383-391.



13. Gram, L., Lovold, T., Melchiorson, J. and Spanggaard, B. 2001. *In vitro* antagonism of Probiotic *Pseudomonas fluorescens* strain AH2 against *Aeromonas salmonicida* does not confer protection of salmon against furunculosis. *Aquacult.* 199: 1-11.
14. Irianto, A. and Austin, B. 2002. Use of probiotics to control furunculosis in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *J. Fish. Dis.* 25: 333-342.
15. Kesarcodi-Watson, A., Kaspar, H., Lategan, M.J. and Gibson, L. 2008. Probiotics in aquaculture: The need, principles and mechanisms of action and screening processes. *Aquacult.* 274: 1-14.
16. Palanivelu, V., Vijayavel, K., Ezhilarasi balasubramanian, S. and Balasubramanian, M.P. 2005. Impact of fertilizer (urea) on oxygen consumption and feeding energetics in the fresh water fish *Oreochromis mossambicus*. *Environ. Toxicol. Pharmacol.* 19: 351-355.
17. Roe, J.H. 1955. The determination of sugar in blood and spinal fluid with anthrone reagent. *J. Biol. Chem.* 212: 335-343.
18. Sugita, H., Miyajima, C. and Deguchi, Y. 1991. The vitamin B12 - producing ability of the intestinal microflora of freshwater fish. *Aquacult.* 92: 267-276.

**Cite this Article as:**

Gopinath, J., Ramanibai, R., Krishnakumar, V. and Inbasekaran, S. 2019. Studies on Biochemical Parameters of Probiotic Supplemented Diet on Rohu (*Labeo rohita*) Fresh Water Fish Fingerlings. *J. Acad. Indus. Res.* 7(12): 172-175.